



The  
Patent  
Office

PCT/GB 98/04031

06 DECEMBER 1999

INVESTOR IN PEOPLE

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

09/857465

GB99/4021

REC'D 26 JAN 2000	
WIPO	PCT


The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

*Andrew Gurney*

Dated 11 January 2000

THIS PAGE BLANK (USPTO,


05 DEC 1998

07DEC98 F409703-1 010007  
551/7700 0.00 - 9826700.8

1/77

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH

1. Your reference	P1339			
2. Patent application number (The Patent Office will fill in this part)	<b>9826700.8</b>			
3. Full name, address and postcode of the or of each applicant (underline all surnames)	UNIVERSITY OF DURHAM SOUTH ROAD DURHAM DH1 3LE			
Patents ADP number (if you know it)				
If the applicant is a corporate body, give the country/state of its incorporation				
4. Title of the invention	PROCESS FOR PREPARING CHIRAL COMPOUNDS			
5. Name of your agent (if you have one)	MARKGRAAF PATENTS LIMITED			
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	The Crescent 54 Blossom Street YORK YO24 1AP			
Patents ADP number (if you know it)				
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	<table border="0"> <tr> <td style="text-align: center;">Country</td> <td style="text-align: center;">Priority application number (if you know it)</td> <td style="text-align: center;">Date of filing (day / month / year)</td> </tr> </table>	Country	Priority application number (if you know it)	Date of filing (day / month / year)
Country	Priority application number (if you know it)	Date of filing (day / month / year)		
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	<table border="0"> <tr> <td style="text-align: center;">Number of earlier application</td> <td style="text-align: center;">Date of filing (day / month / year)</td> </tr> </table>	Number of earlier application	Date of filing (day / month / year)	
Number of earlier application	Date of filing (day / month / year)			
8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. See note (d))	YES			

7083256001

7296486002

9. Enter the number of sheets for any of the following items you are filling with this form. Do not count copies of the same document

Continuation sheets of this form

Description

22

Claim(s)

Abstract

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

*Mark Graaf*

Date 04.12.1998

MARKGRAAF PATENTS LIMITED

12. Name and daytime telephone number of person to contact in the United Kingdom

Philippa M Allen

01904 610586

#### Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

#### Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500303.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.


## PROCESS FOR PREPARING CHIRAL COMPOUNDS

The present invention relates to a process for the preparation of a class of enantiomerically pure chiral compounds, the compounds obtained thereby and novel compounds, compositions thereof and the use thereof as or in the preparation of a pharmaceutical, veterinary product, agrochemical, polymer, library of compounds and their respective intermediates.


Efficient and simple synthesis of known and novel compounds can be the key to commercial success and may also lead to further development and discoveries enabled by availability of compounds in significant purities, yields and the like. Nevertheless development of new synthetic routes is costly and time consuming, without the guarantee of success.

We have now surprisingly found a process for synthesising a class of compounds in novel manner to produce enantiomerically pure hetero compounds.

Accordingly in a first aspect there is provided a process for the preparation of chiral compounds of formula I:



comprising contacting a compound of formula II:



with a source of hydrogen or fluorine;

wherein

A is a chiral centre;

X is selected from oxygen, sulphur and nitrogen;

5 n is selected from 0 and 1 and is equal to the valence of X less 2;

Each  $R^1$  is independently selected from hydrogen, straight chain and branched, saturated and unsaturated  $C_{1-8}$  hydrocarbon optionally substituted by one or more hydroxy, halo, aryl, cyclo  
10  $C_{1-8}$  alkyl and the like;

B is a fragment  $CR^3_2$  wherein each  $R^3$  is independently selected from O-, N- and S- containing functional groups such as hydrogen, halo, azides and cyanides; straight and branched chain,  
15 saturated and unsaturated  $C_{1-4}$  alkyl, alkenyl and alkynyl and aryl, each optionally substituted by hydroxy, halo, saturated or unsaturated  $C_{1-4}$  alkyl, alkenyl or alkynyl, aryl, cyclo  $C_{1-6}$  alkyl, carbonyl, carboxyl, amino, amido, (thio)ether, haloalkyl, silylalkyl and the like;

20 each  $R^2$  is independently selected from hydrogen, straight chain and branched, saturated and unsaturated  $C_{1-8}$  alkyl, optionally substituted by hydroxy, halo, aryl, cyclo  $C_{1-6}$  alkyl, carbonyl, carboxyl, amino, amido, (thio)ether and the like; and

25 one of  $R^1$  and one of  $R^2$  together may form an alkylene group as part of a heterocyclic ring;

with the proviso that when X is nitrogen, n is 2, one of  $R^1$  and two of  $R^2$  are  
30 hydrogen, B is  $CHPh_2$ , the other  $R^1$  and  $R^2$  do not form together a five membered heterocyclic ring.

Preferably X is nitrogen whereby n is 1.

Preferably B is a fragment  $CR^3_2$  wherein  $R^3$  is selected from ethenyl, propenyl ethynyl and propynyl, optionally substituted phenyl, for example 4-methoxy or  
5 4-perfluoryl alkyl phenyl, naphthyl, methyl phenyl and the like.

More preferably B is a group as hereinbefore defined wherein at least one and preferably both of  $R^3$  are aryl, more preferably optionally substituted phenyl.

10 Preferably  $R^2$  is selected from optionally hydroxy, halo, alkoxy substituted branched and straight chain  $C_{1-6}$  alkyl, including methyl i-propyl, i-butyl, t-butyl; and aryl including phenyl and benzyl.

Preferably X is nitrogen wherein n is one and  $R^1$  does not form a cyclic ring  
15 with one of  $R^2$ , i.e. the compound is a non cyclic secondary amine, or  $R^1$  is H, and  $R^2$  is other than H, i.e. the compound is a primary amine.

Contacting the compound of formula II as hereinbefore defined is suitably in the presence of a catalyst which may be homogeneous or heterogeneous, and is  
20 preferably heterogeneous.

Without being limited to this theory it is thought that the conversion according to the process of the invention proceeds via a substitution with subsequent decarboxylation or decarboxylation with subsequent quenching. The catalyst  
25 may be selected from any catalyst suitable for the conversion as hereinbefore defined. Preferably the catalyst comprises a hydrogenation or fluorination catalyst or agent. A hydrogenation catalyst suitably comprises a metal adapted to catalyse a hydrogenation reaction, for example selected from the transition metals of Group VIII of the Periodic Table of the Elements, preferably selected  
30 from Pt, Pd, Ni, Co, Cu, Ru, Fe and Ag and mixtures thereof. The catalyst may be in the form of the metal(s) or salts thereof, optionally in the presence of or

including additional catalytic components or catalytic supports such as C. More preferably the catalyst comprises palladium and carbon, and reaction is in the presence of gaseous hydrogen.

- 5 The catalyst is present in catalytically effective amount.

The process may be carried out with use of solvents, and may be carried out at ambient or elevated temperature and/or pressure, and is preferably carried out at elevated pressure in the range 1-10 atm.

10

- The process of the invention is preferably suited for the preparation of pharmaceutical, veterinary product, agrochemical and polymeric compounds and libraries of such compounds, and their synthetic intermediates. It is a particular advantage of the process of the invention that such compounds may be readily prepared in which B is analogous electronically and/or sterically to characteristic groupings in known pharmaceutical, veterinary product and agrochemicals. The process therefore provides a known route to access compounds and whole ranges of new analogues, wherein the group B is as hereinbefore defined.

20

Alternatively the process as hereinbefore defined is suited for the preparation of metal complexes as asymmetric catalysts.


- In a further aspect of the invention there is provided a class of novel enantiomerically pure chiral hetero compounds of the formula I as hereinbefore defined wherein A, B and  $R^1$  are as hereinbefore defined, X is N and n is 1 with the exception that  $R^2$  is not phenyl or benzyl when  $R^1$  is hydrogen and BH is phenyl or  $CH_3$ .

- 30 Compounds of the formula II as hereinbefore defined may be obtained commercially or prepared by known means. Akiba *et al*, Tetrahedron, 1994, 50




(13), 3905 discloses the preparation of a compound of formula II by cyclisation of amino alcohol with trichloromethyl chloroformate ( $\text{Cl}_3\text{COCOCl}$ ) in the presence of triethylamine ( $\text{Et}_3\text{N}$ ). Using this process compounds of formula II are obtained from compounds of formula III:

5



10 Intermediate compounds of formula III as hereinbefore defined may be obtained commercially or using the process, for example of Gawley and Zhang, J. Org. Chem., 1996, 61, 8103, and Itsuno *et al*, J. Chem. Soc., Perkin Trans. I, 1985, 2039. In these publications is taught the preparation of a compound of formula III as hereinbefore defined by reaction of a compound of formula IV:

15



20 with a compound of formula V:




Reaction is preferably under reflux in cold solvent.

25

Compounds of formula IV and V are commercially available or may be synthesised by known means.

30

In a further aspect of the invention there is provided a process for the preparation of enantiomerically pure chiral polymers comprising a repeating unit of the formula II:



10


wherein P is derived from a polymerisable monomer or oligomer and X,  $R^1$ ,  $R^2$ , B and A are as hereinbefore defined.

Polymerisable monomers may be any known monomers, for example selected from monomers of thermoset and thermoplast polymers and mixtures thereof, including monomers preferably selected from the group consisting of: an epoxy resin such as an epoxy resin derived from the mono or poly-glycidyl derivative of one or more of the group of compounds consisting of aromatic diamines, aromatic monoprimary amines, aminophenols, polyhydric phenols, polyhydric alcohols, polycarboxylic acids and the like; an addition-polymerisation resin, such as a bis-maleimide resin, acrylic, vinyl or unsaturated polyester; a formaldehyde condensate resin, such as a formaldehyde-phenol resin, urea, melamine or phenol resin; a cyanate resin; and an isocyanate resin; polyaromatics such as polysulphones and polyethersulphones; monomers of natural polymers including carbohydrates, polypeptides and proteins including starch, celluloses, collagen, gelatin, dextrans, alginates, chitin and chitosan; and monomers of biodegradable and/or biocompatible polymers such as polyesters including poly(lactic acid), poly(glycolic acid), polycaprolactone and the like, polyorthoesters, polyanhydrides, polyaminoacids and azo polymers, for example for the delivery of a pharmaceutical, veterinary product or agrochemical *in situ*.

In a further aspect of the invention there is provided a process for the preparation of compounds of the formula Iii:

5


(Iii)



10 by the functional modification of a compound of formula I as hereinbefore defined to include additional groups  $R^1$  and  $R^3$  or the interconversion of one compound of formula I as hereinbefore defined to another compound of formula I as hereinbefore defined.

15 Preferably the compound of formula Iii as hereinbefore defined is a spatial, electronic or reactive analogue of a known pharmaceutical, veterinary product, or agrochemical, for example of a neuro active compound, such as the compound orphenadrine of formula:

20



25

for use in treating Parkinson's Disease or of cardiovascular or gastro-intestinal drugs, immunosuppresants, respiratory agents, musculoskeletal and joint disease drugs, immunological products and vaccines, pest control agents, plant growth control agents, plant disease control agents and the like.

In a further aspect of the invention there is provided the use of one or more compounds of formula I as hereinbefore defined in the preparation of a library of compounds comprising:

30

reacting one or more compounds of formula I as hereinbefore defined with one or more substrates which are supported or contained in solid or liquid phase each on an individual support or within an individual vessel; and

5

labelling the support or vessel with means to identify the synthetic history of the supported or contained compound.

The process for preparing a library of compounds may employ any techniques as known in the art of combinatorial chemistry.

10

In a further aspect of the invention there is provided a process for the preparation of a library of compounds of formula I as hereinbefore defined comprising:

15

reacting one or more compounds of formula IV as hereinbefore defined with a plurality of compounds of formula V as hereinbefore defined, and converting via compounds of formula II as hereinbefore defined to compounds of formula I as hereinbefore defined; and

20

optionally labelling the support or vessel with means to identify the synthetic history of the supported or contained compound.

In a further aspect of the invention there is provided a library of compounds of formula I, II or III as hereinbefore defined.

25

Preferably the library of compounds is suitable for any of the hereinbefore defined uses. The library may be provided in the form of a kit of sample boxes for the intended use. The library may contain two or more compounds, for example ten or more compounds, preferably comprises 50-1,000 compounds of

30

any given formula as hereinbefore defined, optionally including synthetic history identification.

In a further aspect of the invention there is provided a pharmaceutical, veterinary product or agrochemical composition comprising a compound of formula I as hereinbefore defined or derivatives thereof together with suitable diluents, adjuvants, carriers and the like.


The invention is now illustrated in non limiting manner with reference to the examples.

## Examples

### 1.1 New Synthetic Approach to Novel Chiral Amines

#### 1.1.1 Synthesis of (S)-2-amino-1,1-diphenyl-3-methyl-1-butanol (2)

The title compound (2) was readily prepared by the addition of L-valine methyl ester hydrochloride (1) to phenylmagnesium bromide, as depicted in Scheme 1, following the modified method described by Gawley<sup>i</sup> and Zhang (1996), and Itsuno<sup>ii</sup> *et al.* (1985).




Scheme 1

Purification over silica gel, gave (2) as a white solid in moderate yield (36 %).

#### 25 1.1.2 Synthesis of (S)-4-isopropyl-5,5-diphenyl-2-oxazolidinone (3)

In the event, the title compound (3) was readily prepared by the cyclisation of aminoalcohol (2) with trichloromethyl chloroformate (Cl<sub>3</sub>COCOC<sub>2</sub>H<sub>5</sub>) in

the presence of triethylamine ( $\text{Et}_3\text{N}$ ), as shown in **Scheme 2**, following the method described by Akiba<sup>iii</sup> *et al.* (1994).




**Scheme 2**

Upon work-up, the solid residue was loaded on to a sintered funnel and then washed with diethyl ether to obtain the title compound (3) as a white solid in good yield (86 %).

### 1.1.3 Synthesis of (S)-2-amino-3-methyl-1,1-diphenylbutane (4)

In the presence of a catalytic amount of palladium on-activated carbon, 2-oxazolidinone (3) was finally submitted to the hydrogenation in a mixture of AcOH and MeOH under 4-5 atm. pressure, as illustrated in **Scheme 3**.




**Scheme 3**

Upon filtration and re-crystallisation from petroleum ether, the title compound (4) was generated as a white solid in good yield (72 %).

### 1.1.4 Synthesis of (S)-2-amino-1,1,3-triphenyl-1-propanol (6)

The title compound (6), following the modified literature methods of Itsuno<sup>ii,iv</sup> *et al.* (1985), Weber<sup>v</sup> *et al.* (1995) and Dammast and Reißig<sup>vi</sup> (1993), was readily prepared by the portionwise addition of L-phenylalanine ethyl ester hydrochloride (5) to phenylmagnesium bromide, as depicted in **Scheme 4**.




Scheme 4

Recrystallisation gave the title compound (6) as a white solid in low yield (9 %).

- 5 [Phillipa, this yield is not representative, but represents what happened on the day, it could clearly be improved and should not be reported]

#### 1.1.5 Synthesis of (S)-4-benzyl-5,5-diphenyl-2-oxazolidinone (7)

In the event, the title compound (7) was readily prepared by the cyclisation of aminoalcohol (6) with trichloromethyl chloroformate ( $\text{Cl}_3\text{COCOC}\text{Cl}$ ) in the presence of triethylamine ( $\text{Et}_3\text{N}$ ), as shown in Scheme 5, following the method described by Akiba<sup>iii</sup> *et al.* (1994).




Scheme 5

- 15 Upon work-up, the solid residue was loaded on to a sintered funnel and then washed with diethyl ether to obtain the title compound (7) as a white solid in excellent yield (97 %).

#### 1.1.6 Synthesis of (S)-2-amino-1,1,3-triphenylpropane (8)

- In the presence of a catalytic amount of palladium on activated carbon, 2-oxazolidinone (7) was finally subjected to the hydrogenation in a mixture of AcOH and MeOH under 4-5 atm. pressure, as illustrated in Scheme 6.
- 20




Scheme 6

Upon filtration and purification over silica gel, eluting with a 3:7 and 4:6 mixture of AcOEt and petrol, the title compound (8) was obtained as a light-brown solid in good yield (71 %).

10

#### 1.1.7 Synthesis of (S)-2-amino-1,1-diphenyl-1-propanol (10)

The title compound (10), following the literature methods of Itsuno<sup>ii</sup> *et al.* (1985), Weber<sup>v</sup> *et al.* (1995) and Dammas<sup>vi</sup> and Reißig<sup>v</sup> (1993), was readily prepared by the portionwise addition of L-alanine methyl ester hydrochloride (9) to phenylmagnesium bromide, as depicted in Scheme 7.




Scheme 7

Flash column chromatography, eluting with dichloromethane and then further elution with a mixture of AcOEt and petrol, ranging from 15 % up to 100 %, gave the title compound (10) as a white solid in moderate yield (52 %).

#### 1.1.8 Synthesis of (S)-4-methyl-5,5-diphenyl-2-oxazolidinone (11)

In the event, the title compound (11) was readily prepared by the cyclisation of aminoalcohol (10) with trichloromethyl chloroformate (Cl<sub>3</sub>COCOC<sub>2</sub>H<sub>5</sub>) in the presence of triethylamine (Et<sub>3</sub>N), as shown in Scheme 8, following the method described by Akiba<sup>iii</sup> *et al.* (1994).






Scheme 8

5 Upon work-up, the solid residue was loaded on to a sintered funnel and then washed with diethyl ether to obtain the title compound (11) as a white solid in good yield (76 %).

### 1.1.9 Synthesis of (S)-2-amino-1,1-diphenyl-propane (12)

10 In the presence of a catalytic amount of palladium on activated carbon, 2-oxazolidinone (11) was finally subjected to the hydrogenation in a mixture of AcOH and MeOH under 4-5 atm. pressure, as illustrated in Scheme 9.




Scheme 9

15 Upon filtration and purification by dry-flash column chromatography, eluting first with AcOEt, and then with a mixture of MeOH and AcOEt, ranging from 5 % up to 30 %, gave the title compound (12) as a white solid in moderate yield (71 %).

## 1.2 Experimental

### 1.2.1 (S)-2-amino-1,1-diphenyl-3-methyl-1-butanol (2)




THIS PAGE BLANK (USPTO)

## (2)

L-Valine methyl ester hydrochloride (9.9 g, 59.06 mmol) was added portionwise to a 1.0 M solution of phenylmagnesium bromide (108.8 g, 0.6 mol) in THF at 0 °C and heated at reflux for 20h. After quenching with crushed ice and NH<sub>4</sub>Cl salt, the organic layer was separated, washed with brine and concentrated under reduced pressure. The resulting solid was treated with HCl (2.0 M, 100 ml) and then evaporated to dryness under reduced pressure. Impurities precipitated out as a white solid, when the amine hydrochloride salt was dissolved in hot MeOH and allowed to cool to room temperature. After removing the impurities by filtration, the filtrate was made basic with KOH (1.0 M) and the organics were extracted into diethyl ether (4x100 ml). Combined organic extracts were dried over MgSO<sub>4</sub> and concentrated under reduced pressure to obtain a crude product as a light brown solid. Purification over silica gel, eluting with a 1:4 and 1:1 mixture of ethyl acetate and petrol gave the title compound (2) (5.42 g, 36 %) as a white solid. m.p. 90-92 °C (lit<sup>i</sup> 94-95 °C).  $[\alpha]_D^{25} = -107.92^\circ$  (c, 0.0424 in CHCl<sub>3</sub>) (lit<sup>ii</sup>: -127.7° (c, 0.639 in CHCl<sub>3</sub>).  $\delta_H$  0.81 (3H, d, <sup>3</sup>J= 6.90 Hz, CH<sub>3</sub>), 0.85 (3H, d, <sup>3</sup>J= 7.20 Hz), 1.67 (1H, ds, <sup>3</sup>J= 1.80 and 6.90 Hz, CH-Me<sub>2</sub>), 3.76 (1H, d, <sup>3</sup>J= 2.10 Hz, CH-NH<sub>2</sub>), 7.04-7.58 (10H, m, Ar).  $\delta_C$  16.3 and 23.2 (CH<sub>3</sub>), 28.1 (CH-Me<sub>2</sub>), 60.4 (CH-NH<sub>2</sub>), 79.9 (C-OH), 125.7, 126.1, 126.5, 126.8, 128.2 and 128.6 (o-, m- and p-Ar), 145.1 and 148.2 (α-Ar). Anal. Calcd. for C<sub>17</sub>H<sub>21</sub>NO: C 79.96; H 8.29; N 5.48. Found: C 79.80; H 8.15; N 5.39. ir 3338 (OH and NH<sub>2</sub>). m/e (CI-CH<sub>4</sub>) 256 (MH<sup>+</sup>, 14 %), 72 (100 %).

## 1.2.2

## (S)-4-isopropyl-5,5-diphenyl-2-oxazolidinone (3)




5 Trichloromethyl chloroformate (2.71 g, 13.7 mmol) was added to a mixture of (S)-2-amino-3-methyl-1,1-diphenyl-1-butanol (**2**) (3.18 g, 12.45 mmol) and triethylamine (2.68 g, 26.52 mmol) in  $\text{CH}_2\text{Cl}_2$  at 0 °C. The reaction mixture was stirred for 2h at the same temperature and then poured into a brine solution (250 ml). The aqueous layer was made basic with NaOH pellets and organic products were

10 extracted into AcOEt (5x200 ml). Combined organic extracts were dried over  $\text{MgSO}_4$  and concentrated under reduced pressure. The resulting crude product was washed with diethyl ether to obtain the title compound (**3**) (3.03 g, 86%) as a white solid. m.p. 250-251 °C (lit. 250-251 °C).  $[\alpha]_{\text{D}}^{25} = -201.59^\circ$  (c, 0.0252 in DMSO).  $\delta_{\text{H}}$  (DMSO- $\text{d}_6$ ) 0.51 (3H, d,  $^3J = 6.60$  Hz,  $\text{CH}_3$ ), 0.92 (3H, d,  $^3J = 7.20$  Hz,  $\text{CH}_3$ ), 1.86 (1H, ds,  $^3J = 2.10$  and 6.60 Hz,  $\text{CH-Me}_2$ ), 4.46 (1H, d,  $^3J = 6.5$  Hz,  $\text{CH-NH}_2$ ), 7.24-7.72 (10H, m, Ar-H), 8.14 (1H, s, NH).  $\delta_{\text{C}}$  15.2 and 20.9 ( $\text{CH}_3$ ), 29.8 (CH), 64.9 (CH-NHCO), 88.4 (C-O), 125.8, 126.2, 127.9, 128.4, 128.8 and 129.1 (Ar), 140.5 and 146.1 ( $\alpha$ -Ar), 158.1 (C=O). Ir 3295 ( $\text{NH}_2$ ), 1765 and 1745 (C=O). m/e (CI- $\text{NH}_3$ ) 282 ( $\text{MH}^+$ , 25 %), 299 ( $\text{MNH}_4^+$ , 8 %), 238 (96 %), 223 (100 %), 72 (100 %).

15

20

1.2.3

**(S)-2-amino-3-methyl-1,1-diphenylbutane (4)**

5 A solution of (S)-4-isopropyl-5,5-diphenyl-2-oxazolidinone (3) (2.9 g, 10.31 mmol) in MeOH/AcOH and a 10 % Pd (435 mg, 4.09 mmol) on activated carbon was shaken for 68h under 4-5 atm pressure of hydrogen at room temperature. The catalyst was filtered off over Hyflo Super Cell and organic solvents were evaporated under reduced pressure. The resulting residue was treated with HCl (2.0


10 M, 50 ml), stirred for 2h at room temperature, made basic with NaOH pellets, and saturated with K<sub>2</sub>CO<sub>3</sub> and NaCl. Organic compounds were then extracted into AcOEt (3x 100 ml), dried over MgSO<sub>4</sub>/K<sub>2</sub>CO<sub>3</sub> and concentrated under reduced pressure to obtain a crude product. Re-crystallisation from petroleum ether gave the title compound (4) (1.79 g, 72 %) as a light-brown solid. m.p. 71-72 °C.  $[\alpha]_D^{25} = -4.19^\circ$  (c,

15 0.1097 in CHCl<sub>3</sub>).  $\delta_H$  0.78 (3H, d,  $^3J = 6.60$  Hz, CH<sub>3</sub>), 0.91 (3H, d,  $^3J = 7.20$  Hz, CH<sub>3</sub>), 1.26 (2H, broad s, NH<sub>2</sub>), 1.62 (1H, ds, CHMe<sub>2</sub>), 3.45 (1H, dd,  $^3J = 10.5$  and 2.40 Hz, CH-NH<sub>2</sub>), 3.70 (1H, d,  $^3J = 10.5$  Hz, CH-Ph<sub>2</sub>), 7.00-7.40 (10H, m, Ar-H).  $\delta_C$  14.2 and 21.5 (CH<sub>3</sub>), 28.9 (CH-Me<sub>2</sub>), 58.1 and 58.9 (CH-NH<sub>2</sub> and CH-Ph<sub>2</sub>), 126.5, 126.7, 128.2, 128.5, 128.8 and 129.0 (o-, m- and p-Ar), 143.5 (2x $\alpha$ -Ar). Anal. Calcd for C<sub>17</sub>H<sub>21</sub>N: C 85.30; H 8.84; N 5.85. Found: C 85.12; H 8.91; N 5.96. ir 3361 (NH<sub>2</sub>).

20 m/e (CI-CH<sub>4</sub>) 240 (MH<sup>+</sup>, 8 %), 72 (100 %).

1.2.4


## (S)-2-Amino-1,1,3-triphenyl-1-propanol (6)



- 5 L-Phenylalanine ethyl ester hydrochloride (9.9 g, 43.1 mmol) was added portionwise to a 1.0 M solution of phenylmagnesium bromide (63.46 g, 0.35 mol) in THF at 0 °C and stirred for 20h at room temperature. After quenching with crushed ice and concentrated HCl, the aqueous layer was separated and evaporated to dryness under reduced pressure. The resulting solid was washed with diethyl ether and AcOEt
- 10 to obtain a white gummy HCl-salt. Upon basification with NaOH (1.0 M), organic products were extracted into diethyl ether and AcOEt, dried over MgSO<sub>4</sub>, and concentrated under reduced pressure to obtain a crude product. Re-crystallisation from a mixture of AcOEt and diethyl ether gave the title compound (6) (1.16 g, 9 %) as a white solid. m.p. 141-142 °C (lit<sup>ii</sup> 144-145 °C; lit<sup>vi</sup> 143-144 °C).  $[\alpha]_D^{25} = -88.40^\circ$
- 15 (c, 0.0181 in CHCl<sub>3</sub>) (lit<sup>ii</sup> - 88.50° (c, 0.604 in CHCl<sub>3</sub>); lit<sup>vi</sup>: - 94.3° (c, 2.30 in CHCl<sub>3</sub>).  $\delta_H$  2.38 (1H, dd,  $^3J = 10.8$  Hz,  $^2J = 13.8$  Hz, CH<sub>2</sub>-Ph), 2.58 (1H, dd,  $^3J = 2.4$  Hz,  $^2J = 13.8$  Hz, CH<sub>2</sub>-Ph), 4.11 (1H, dd,  $^3J = 2.4$  Hz,  $^3J = 10.8$  Hz, CH-NH<sub>2</sub>), 7.06-7.62 (15H, m, Ar-H).  $\delta_C$  36.9 (CH<sub>2</sub>-Ph), 58.4 (CH-NH<sub>2</sub>), 78.7 (C-OH), 125.6, 126.0, 126.6, 126.7, 126.9, 128.4, 128.7, 128.8 and 129.3 (o-, m- and p-Ar), 139.8, 144.5 and 147.0 ( $\alpha$ -Ar). ir 3365 (NH<sub>2</sub>), 3320 (OH). m/e (CI-NH<sub>3</sub>) 304 (MH<sup>+</sup>, 30 %), 271 (100 %).
- 20

1.2.5


## (S)-4-benzyl-5,5-diphenyl-2-oxazolidinone (7)



5 Trichloromethyl chloroformate (718 mg, 3.63 mmol) was added to a mixture of (S)-2-amino-1,1,3-triphenyl-1-propanol (6) (1.00 g, 3.30 mmol) and triethylamine (710 mg, 7.02 mmol) in  $\text{CH}_2\text{Cl}_2$  at 0 °C. The reaction mixture was stirred for 5h at the same temperature and then poured into a brine solution (150 ml). The aqueous layer was made basic with powdered  $\text{K}_2\text{CO}_3$  and organics were extracted

10 into dichloromethane (3x50 ml). The combined organic extracts were dried over  $\text{MgSO}_4/\text{K}_2\text{CO}_3$  and concentrated under reduced pressure. The resulting crude product was washed with diethyl ether to obtain the title compound (7) (1.06 g, 97 %) as a white solid. m.p. 259-261 °C (lit ? °C).  $[\alpha]_D^{25} = -241.94^\circ$  (c, 0.0211 in DMSO),  $\delta_{\text{H}}$  (DMSO- $d_6$ ) 2.18 (1H, dd,  $^3J = 10.8$  Hz,  $^2J = 13.8$  Hz,  $\text{CH}_2\text{-Ph}$ ), 2.52 (1H, dd,  $^3J = 3.6$  Hz,  $^2J = 13.8$  Hz,  $\text{CH}_2\text{-Ph}$ ), 4.67 (1H, dd,  $^3J = 3.6$  Hz,  $^3J = 10.8$  Hz,  $\text{CH-NH}_2$ ), 6.90-7.60 (15H, m, Ar-H).  $\delta_{\text{C}}$  44.2 ( $\text{CH}_2\text{-Ph}$ ), 50.5 ( $\text{CH-NH}$ ), 94.1 (C-O), 130.5, 130.9, 131.5, 132.6, 132.8, 133.0, 133.1, 133.3 and 133.4 (o-, m- and p-Ar), 141.1, 143.4 and 146.5 ( $\alpha\text{-Ar}$ ), 163.7 (C=O). ir 3248 ( $\text{NH}_2$ ), 1760 and 1725 (C=O). m/e (CI- $\text{NH}_3$ ) 330 ( $\text{MH}^+$ , 5 %), 347 ( $\text{MNH}_4^+$ , 6 %), 196 (100 %).

## 1.2.6 (S)-2-Amino-1,1,3-triphenyl-propane (8)



5           A solution of (S)-4-benzyl-5,5-diphenyl-2-oxazolidinone (7) (940 mg, 2.85 mmol) in MeOH/AcOH and a 10 % Pd (121 mg, 1.14 mmol) on activated carbon was shaken for 43h under 4-5 atm pressure of hydrogen at room temperature. The catalyst was filtered off over Hyflo Super Cell and organic solvents were evaporated under reduced pressure. The resulting residue was treated with HCl, stirred for 2h at


10 room temperature, made basic with NaOH pellets, and saturated with  $K_2CO_3$  and NaCl. Organics were then extracted into dichloromethane (4x 50 ml), dried over  $MgSO_4/K_2CO_3$  and concentrated under reduced pressure to obtain a crude product. Purification over silica gel, eluting with a 3:7 and 4:6 mixture of AcOEt and petroleum ether, gave the title compound (8) (584 mg, 71 %) as a light-brown solid.

15 m.p. 71-72 °C.  $[\alpha]_D^{25} = -8.03^\circ$  (c, 0.1046 in  $CHCl_3$ ).  $\delta_H$  1.21 (2H, broad s,  $NH_2$ ), 2.29 (1H, dd,  $^3J = 9.6$  Hz,  $^2J = 13.5$  Hz,  $CH_2$ -Ph), 2.79 (1H, dd,  $^3J = 2.1$  Hz,  $^2J = 13.2$  Hz,  $CH_2$ -Ph), 3.71 (1H, d,  $^3J = 9.9$  Hz, CH- $Ph_2$ ), 3.81 (1H, ddd,  $^3J = 2.7, 9.9$  and 12.6 Hz, CH- $NH_2$ ), 7.06-7.33 (15H, m, Ar-H).  $\delta_C$  41.9 ( $CH_2$ -Ph), 55.7 and 59.7 (CH- $Ph_2$  and CH- $NH_2$ ), 126.3, 126.5, 126.6, 128.1, 128.2, 128.4, 128.7, 128.8 and 129.1 (o-, m- and p-Ar), 139.7, 142.6 and 143.1 ( $\alpha$ -Ar). ir 3387 ( $NH_2$ ). m/e (CI- $NH_3$ ) 288 ( $MH^+$ , 100 %).

20




## 1.2.7 (S)-2-Amino-1,1-diphenyl-1-propanol (10)



- 5 L-Alanine methyl ester hydrochloride (9.9 g, 70.9 mmol) was added portionwise to a 1.0 M solution of phenylmagnesium bromide (78.0 g, 0.43mol) in THF at 0 °C and then heated under reflux for 21h. The reaction mixture was cooled to 0 °C, quenched with dropwise addition of saturated NH<sub>4</sub>Cl, and stirred for 1h. After collecting insoluble products through the Buchner funnel, organic products were
- 10 extracted into AcOEt (3x100 ml). The combined organic extracts were dried over K<sub>2</sub>CO<sub>3</sub>/MgSO<sub>4</sub>, and concentrated under reduced pressure to obtain a crude product. Impurities were washed with dichloromethane over silica gel by means of dry-flash column chromatography, further elution with a mixture of AcOEt and petrol, ranging from 20 % up to 100 %, gave the title compound (10) (1.16 g, 9 %) as a white solid.
- 15 m.p. 100-101 °C (lit<sup>ii,v</sup> 100-102 °C).  $[\alpha]_D^{25} = -85.59^\circ$  (c, 0.0362 in CHCl<sub>3</sub>) (lit<sup>ii</sup>: -82.38° (c, 0.814 in CHCl<sub>3</sub>; lit<sup>v</sup>: -85.9° (c, 2.77 in CHCl<sub>3</sub>).  $\delta_H$  0.94 (3H, d, <sup>3</sup>J= 6.30 Hz, CH<sub>3</sub>), 1.23 (2H, broad s, NH<sub>2</sub>), 4.15 (1H, q, <sup>3</sup>J= 6.30 Hz, CH-NH<sub>2</sub>), 4.25 (1H, broad s, OH), 7.10-7.66 (10H, m, Ar-H).  $\delta_C$  17.4 (CH<sub>3</sub>), 52.1 (CH-NH<sub>2</sub>), 78.7 (C-OH), 125.7, 126.1, 126.6, 126.9, 128.2 and 128.7 (o-, m- and p-Ar), 145.0 and 147.2 ( $\alpha$ -Ar). Anal.
- 20 Calcd. for C<sub>15</sub>H<sub>17</sub>NO: C 79.26; H 7.54; N 6.16. Found: C 79.30; H 7.66; N 6.27. ir 3432 (OH), 3389 (NH<sub>2</sub>). m/e (CI-NH<sub>3</sub>) 228 (MH<sup>+</sup>, 100 %).

1.2.8

**(S)-4-Methyl-5,5-diphenyl-2-oxazolidinone (11)**


5 Trichloromethyl chloroformate (6.37 g, 32.19 mmol) was added to a mixture of (S)-2-amino-1,1-diphenyl-1-propanol (10) (6.65 g, 29.26 mmol) and triethylamine (6.31 g, 62.3 mmol) in  $\text{CH}_2\text{Cl}_2$  at 0 °C. The reaction mixture was stirred for 5h at the same temperature, poured into a brine solution (150 ml), and diluted with more dichloromethane. After collecting insoluble impurities through the Buchner

10 funnel, the organic layer was separated and the aqueous layer was washed once with a mixture of dichloromethane and AcOEt. The combined organic extracts were dried over  $\text{MgSO}_4/\text{K}_2\text{CO}_3$  and concentrated under reduced pressure. The resulting crude product was washed with diethyl ether, water, AcOEt and diethyl ether again, to obtain the title compound (11) (5.67 g, 76 %) as a white solid. m.p. 264-266 °C  $[\alpha]_D^{25}$

15 = -279.71° (c, 0.0414 in DMSO).  $\delta_{\text{H}}$  0.82 (3H, d,  $^3J = 6.30$  Hz,  $\text{CH}_3$ ), 4.65 (1H, q,  $^3J = 6.0$  Hz, CH-NH<sub>2</sub>), 7.10-7.70 (10H, m, Ar-H), 7.93 (1H, broad s, NH).  $\delta_{\text{C}}$  19.6 ( $\text{CH}_3$ ), 55.9 (CH-NH<sub>2</sub>), 85.6 (C-O), 126.3, 126.4, 128.1, 128.6, 128.8 and 129.1 (o-, m- and p-Ar), 140.6 and 144.2 ( $\alpha$ -Ar), 157.6 (C=O). ir 3254(NH), 1745 and 1725 (C=O). m/e (CI-NH<sub>3</sub>) 254 ( $\text{MH}^+$ , 9 %), 271 ( $\text{MNH}_4^+$ , 55 %), 52 (100 %).

20

## 1.2.9 (S)-2-Amino-1,1-diphenyl-propane (12)



5 A suspension of (S)-4-methyl-5,5-diphenyl-2-oxazolidinone (11) (3.52 g, 13.90 mmol) in MeOH/AcOH and a 10 % Pd (148 mg, 1.39 mmol) on activated carbon was shaken for 45h under 4-5 atm pressure of hydrogen at room temperature. The catalyst was filtered off over Hyflo Super Cell and organic solvents were evaporated under reduced pressure. The resulting residue was treated with HCl (2M, 100 ml), stirred overnight at room temperature, made basic with NaOH pellets, and saturated with K<sub>2</sub>CO<sub>3</sub>. The organics were then extracted into diethyl ether (3x 100 ml), dried over MgSO<sub>4</sub>/K<sub>2</sub>CO<sub>3</sub> and concentrated under reduced pressure to obtain a crude product. Impurities were washed with AcOEt over silica gel by means of dry-  
 10 flash column chromatography, and then further elution with a mixture of MeOH and AcOEt, ranging from 5 % up to 30 %, gave the title compound (12) (1.90 g, 65 %) as a white solid. m.p. 76-77 °C.  $[\alpha]_D^{25} = -19.32$  (c, 0.10765 in CHCl<sub>3</sub>).  $\delta_H$  1.04 (3H, d, <sup>3</sup>J= 6.30 Hz, CH<sub>3</sub>), 1.31 (2H, broad s, NH<sub>2</sub>), 3.55 (1H, d, J= 9.90 Hz, CH-Ph<sub>2</sub>), 3.73 (1H, dq, <sup>3</sup>J= 6.30 and 10.20 Hz, CH-NH<sub>2</sub>), 7.10-7.40 (10H, m, Ar-H).  $\delta_C$  22.4 (CH<sub>3</sub>), 50.3 (CH-NH<sub>2</sub>), 62.4 (CH-Ph<sub>2</sub>), 126.5, 126.8, 128.2, 128.5, 128.7 and 129.0 (o-, m- and p-Ar), 143.3 and 143.7 ( $\alpha$ -Ar). Anal. Calcd for C<sub>15</sub>H<sub>17</sub>NO: C 85.26; H 8.11; N 6.63. Found: C 85.10; H 8.08; N 6.36. ir 3343 (NH<sub>2</sub>). m/e (CI-NH<sub>3</sub>) 212 (MH<sup>+</sup>, 100 %).

25

30

**THIS PAGE BLANK (USPTO)**